

Anti CYP2B6 Antibody

Catalog No: tcsa13515



Available Sizes

Size: 100μl



Specifications

Application:

WB, IHC

Species Reactivity:

Human, Mouse, Rat

Host Species:

Rabbit

Immunogen / Amino acids:

Recombinant protein of human CYP2B6

Conjugation:

Unconjugated

Clonality:

Polyclonal

Isotype:

IgG

Form:

Liquid

Storage Buffer:

PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Recommended Dilution:

WB 1:500 - 1:1000

IHC 1:50 - 1:100

Storage Instruction:

Store at -20C. Avoid freeze / thaw cycles.

Alternative Names:


CYP2B6 antibody; Cytochrome P450 2B6 antibody; CYP2B6 antibody

SwissProt:


P20813

Product Description


This gene, CYP2B6, encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by phenobarbital. The enzyme is known to metabolize some xenobiotics, such as the anti-cancer drugs cyclophosphamide and ifosphamide. Transcript variants for this gene have been described; however, it has not been resolved whether these transcripts are in fact produced by this gene or by a closely related pseudogene, CYP2B7. Both the gene and the pseudogene are located in the middle of a CYP2A pseudogene found in a large cluster of cytochrome P450 genes from the CYP2A, CYP2B and CYP2F subfamilies on chromosome 19q.




Western blot analysis of extracts of various cell lines, using CYP2B6 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 90s.




Immunohistochemistry of paraffin-embedded mouse brain using CYP2B6 antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded mouse kidney using CYP2B6 antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human lung using CYP2B6 antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human colon using CYP2B6 antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human colon carcinoma using CYP2B6 antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human breast cancer using CYP2B6 antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human stomach using CYP2B6 antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded human gastric cancer using CYP2B6 antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded mouse lung using CYP2B6 antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded mouse liver using CYP2B6 antibody at dilution of 1:100 (40x lens).

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!