



Anti VTN Antibody

Catalog No: tcsa15257

| Available Sizes |
|--|
| ze: 100μl |
| Specifications |
| oplication: B, IHC, IF |
| pecies Reactivity: uman,Mouse,Rat |
| ost Species: abbit |
| nmunogen / Amino acids: ecombinant protein of human VTN |
| onjugation: nconjugated |
| onality: onoclonal |
| otype: G |
| orm: quid |

Storage Buffer:

PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Recommended Dilution:

WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200



Storage Instruction:

Store at -20C. Avoid freeze / thaw cycles.

Alternative Names:


VTN antibody; Vitronectin antibody; VN antibody; S-protein antibody; Serum-spreading factor antibody; V75 antibody; VTN antibody


SwissProt:


P04004

Product Description


The protein encoded by this gene is a member of the pexin family. It is found in serum and tissues and promotes cell adhesion and spreading, inhibits the membrane-damaging effect of the terminal cytolytic complement pathway, and binds to several serpin serine protease inhibitors. It is a secreted protein and exists in either a single chain form or a clipped, two chain form held together by a disulfide bond.








Immunohistochemistry of paraffin-embedded human liver using VTN antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human liver cancer using VTN antibody at dilution of 1:100 (40x lens).







Immunohistochemistry of paraffin-embedded mouse skin using VTN antibody at dilution of 1:100 (40x lens).




Immunofluorescence analysis of HepG2 cells using VTN antibody.



Immunofluorescence analysis of MCF-7 cells using VTN antibody.



Immunofluorescence analysis of A431 cells using VTN antibody.



Immunofluorescence analysis of A549 cells using VTN antibody.

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!