

# Anti DiMethyl Histone Antibody

Catalog No: tcsa17018



## Available Sizes

---

**Size:** 100μl



## Specifications

---

**Application:**

WB, IHC, IF, IP, CHIP

**Species Reactivity:**

Human, Mouse, Rat, Other (Wide Range)

**Host Species:**

Rabbit

**Immunogen / Amino acids:**

A synthetic methylated peptide corresponding to residues surrounding K27 of human histone H3

**Conjugation:**

Unconjugated

**Isotype:**

IgG

**Form:**

Liquid

**Storage Buffer:**

PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Recommended Dilution:**

WB 1:500 - 1:2000

IHC 1:50 - 1:200

IF 1:50 - 1:200

IP 1:50 - 1:200

ChIP 1:20 - 1:100

CHIPseq 1:20 - 1:100

**Storage Instruction:**

Store at -20C. Avoid freeze / thaw cycles.

**Alternative Names:**


HIST3H3 antibody; Histone H3.1t antibody; H3/t antibody; H3t antibody; H3/g antibody; HIST3H3 antibody; H3FT antibody

**SwissProt:**


Q16695

**Product Description**


Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.




Western blot analysis of extracts of various cell lines, using DiMethyl-Histone H3-K27 antibody. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST.




Immunohistochemistry of paraffin-embedded Rat brain using DiMethyl-Histone H3-K27 antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded Rat kidney using DiMethyl-Histone H3-K27 antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded Rat liver using DiMethyl-Histone H3-K27 antibody at dilution of 1:100 (40x lens).




Immunofluorescence analysis of 293T cells using DiMethyl-Histone H3-K27 antibody. Blue: DAPI for nuclear staining.



Chromatin immunoprecipitation analysis extracts of 293 cell line, using DiMethyl-Histone H3-K27 antibody and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.




Dot-blot analysis of all sorts of methylation peptides using DiMethyl-Histone H3-K27 antibody.




Immunohistochemistry of paraffin-embedded rat lung using DiMethyl-Histone H3-K27 antibody at dilution of 1:200 (40x lens).







Immunohistochemistry of paraffin-embedded human kidney using DiMethyl-Histone H3-K27 antibody at dilution of 1:200 (40x lens).




Immunohistochemistry of paraffin-embedded human lung cancer using DiMethyl-Histone H3-K27 antibody at dilution of 1:200 (40x lens).




Immunohistochemistry of paraffin-embedded human liver cancer using DiMethyl-Histone H3-K27 antibody at dilution of 1:200 (40x lens).



Immunohistochemistry of paraffin-embedded Mouse heart using DiMethyl-Histone H3-K27 antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded Mouse kidney using DiMethyl-Histone H3-K27 antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded Mouse lung using DiMethyl-Histone H3-K27 antibody at dilution of 1:100 (40x lens).

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!