

Anti HMGB1 Antibody

Catalog No: tcsa17048



Available Sizes

Size: 100µl



Specifications

Application:

WB, IHC, IF

Species Reactivity:

Human, Mouse, Rat

Host Species:

Rabbit

Immunogen / Amino acids:

A synthetic peptide of human HMGB1

Conjugation:

Unconjugated

Clonality:

Polyclonal

Isotype:

IgG

Form:

Liquid

Storage Buffer:

PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Recommended Dilution:

WB 1:500 - 1:2000

IHC 1:50 - 1:200

IF 1:50 - 1:200

Storage Instruction:

Store at -20C. Avoid freeze / thaw cycles.

Alternative Names:


HMGB1 antibody; High mobility group protein B1 antibody; High mobility group protein 1 antibody; HMG-1 antibody;
HMGB1 antibody; HMG1 antibody

SwissProt:


P09429

Product Description


This gene encodes a protein that belongs to the High Mobility Group-box superfamily. The encoded non-histone, nuclear DNA-binding protein regulates transcription, and is involved in organization of DNA. This protein plays a role in several cellular processes, including inflammation, cell differentiation and tumor cell migration. Multiple pseudogenes of this gene have been identified. Alternative splicing results in multiple transcript variants that encode the same protein.




Western blot analysis of extracts of various cell lines, using HMGB1 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.




Immunohistochemistry of paraffin-embedded human gastric cancer using HMGB1 Antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human leiomyoma of uterus using HMGB1 Antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human adenomyosis using HMGB1 Antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded mouse lung using HMGB1 Antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded mouse liver using HMGB1 Antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded mouse brain using HMGB1 Antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded mouse heart using HMGB1 Antibody at dilution of 1:100 (40x lens).




Immunofluorescence analysis of HeLa cells using HMGB1 antibody at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.




Immunohistochemistry of paraffin-embedded human amygdalitis using HMGB1 Antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human stomach using HMGB1 Antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human breast cancer using HMGB1 Antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human adenomyosis using HMGB1 Antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded rat lung using HMGB1 Antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded rat brain using HMGB1 Antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded human breast cancer using HMGB1 Antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded human stomach using HMGB1
Antibody at dilution of 1:100 (40x lens).

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!