



# **Anti HNRNPA2B1 Antibody**

**Catalog No: tcsa17065** 

WB 1:500 - 1:2000

IHC 1:50 - 1:200

IF 1:50 - 1:200

A A	vailable Sizes
<b>Size:</b> 100	0μΙ
S <sub>I</sub>	pecifications
<b>Applicat</b> WB, IHC,	
	<b>Reactivity:</b> Iouse,Rat
<b>Host Sp</b> e Rabbit	ecies:
	gen / Amino acids: nant protein of human HNRNPA2B1
<b>Conjuga</b> Unconjug	
<b>Clonality</b> Polyclona	
<b>Isotype:</b> IgG	
Form: Liquid	
<b>Storage</b> PBS with	Buffer: 0.02% sodium azide, 50% glycerol, pH7.3.
Recomm	nended Dilution:





### **Storage Instruction:**

Store at -20C. Avoid freeze / thaw cycles.

#### **Alternative Names:**

HNRNPA2B1 antibody; Heterogeneous nuclear ribonucleoproteins A2/B1 antibody; hnRNP A2/B1 antibody;


HNRNPA2B1 antibody; HNRPA2B1 antibody

#### **SwissProt:**

P22626

## **Product Description**

This gene belongs to the A/B subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins and they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with premRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene has two repeats of quasi-RRM domains that bind to RNAs. This gene has been described to generate two alternatively spliced transcript variants which encode different isoforms.




Nestern blot analysis of extracts of various cell lines, using HNRNPA2B1 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST.




Immunohistochemistry of paraffin-embedded human colon using HNRNPA2B1 Antibody at dilution of 1:200 (40x lens).







Immunohistochemistry of paraffin-embedded human kidney cancer using HNRNPA2B1 Antibody at dilution of 1:200 (40x lens).




Immunohistochemistry of paraffin-embedded human gastric cancer using HNRNPA2B1 Antibody at dilution of 1:200 (40x lens).




Immunofluorescence analysis of HeLa cells using HNRNPA2B1 antibody. Blue: DAPI for nuclear staining.



Immunohistochemistry of paraffin-embedded human colon using HNRNPA2B1 Antibody at dilution of 1:200 (40x lens).





Immunohistochemistry of paraffin-embedded human kidney using HNRNPA2B1 Antibody at dilution of 1:200 (40x lens).




Immunohistochemistry of paraffin-embedded human stomach using HNRNPA2B1 Antibody at dilution of 1:200 (40x lens).








Immunohistochemistry of paraffin-embedded mouse kidney using HNRNPA2B1 Antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded rat brain using HNRNPA2B1 Antibody at dilution of 1:100 (40x lens).

Antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded rat kidney using HNRNPA2B1 Antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded rat liver using HNRNPA2B1 Antibody at dilution of 1:100 (40x lens).

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!