

# Anti AIMP1 Antibody

Catalog No: tcsa18869



**Available Sizes** 

**Size:** 100µl

Specifications

#### **Application:**

WB, IHC, IF

Species Reactivity:

Human,Mouse,Rat

#### **Host Species:**

Rabbit

## Immunogen / Amino acids:

Recombinant protein of human AIMP1

### **Conjugation:**

Unconjugated

## **Clonality:**

Polyclonal

Isotype	
lgG	

## Form: Liquid

#### **Storage Buffer:**

PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

#### **Recommended Dilution:**

WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200



#### **Storage Instruction:**

Store at -20C. Avoid freeze / thaw cycles.

#### **Alternative Names:**


AIMP1 antibody; Aminoacyl tRNA synthase complex-interacting multifunctional protein 1 antibody; Multisynthase complex auxiliary component p43 antibody; AIMP1 antibody; EMAP2 antibody; SCYE1 antibody


#### SwissProt:

Q12904

## **Product Description**

The protein encoded by this gene is a cytokine that is specifically induced by apoptosis, and it is involved in the control of angiogenesis, inflammation, and wound healing. The release of this cytokine renders the tumor-associated vasculature sensitive to tumor necrosis factor. The precursor protein is identical to the p43 subunit, which is associated with the multi-tRNA synthetase complex, and it modulates aminoacylation activity of tRNA synthetase in normal cells. This protein is also involved in the stimulation of inflammatory responses after proteolytic cleavage in tumor cells. Multiple transcript variants encoding different isoforms have been found for this gene. A pseudogene has been identified on chromosome 20.






Western blot analysis of extracts of various cell lines, using AIMP1 Antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000

Immunohistochemistry of paraffin-embedded rat spleen using AIMP1 antibody at dilution of 1:100 (40x lens).


dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.




Web: www.taiclone.com Tel: +886-2-2735-9682 Email: order@taiclone.com




Immunohistochemistry of paraffin-embedded rat kidney using AIMP1 antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded rat heart using AIMP1 antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human gastric cancer using AIMP1 antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded human liver cancer using AIMP1 antibody at dilution of 1:100 (40x lens).






Immunohistochemistry of paraffin-embedded mouse kidney using AIMP1 antibody at dilution of 1:100 (40x lens).

Immunohistochemistry of paraffin-embedded mouse heart using AIMP1 antibody at dilution of 1:100 (40x lens).





Immunofluorescence analysis of HeLa cells using AIMP1 antibody. Blue: DAPI for nuclear staining.

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!