

Anti MCM4 Antibody

Catalog No: tcsa23118



Available Sizes

Size: 100μl



Specifications

Application:

WB

Species Reactivity:

Human

Host Species:

Rabbit

Immunogen / Amino acids:

A synthetic peptide of human MCM4

Conjugation:

Unconjugated

Clonality:

Polyclonal

Isotype:

IgG

Form:

Liquid

Storage Buffer:

PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Recommended Dilution:

WB 1:500 - 1:2000

Storage Instruction:

Store at -20C. Avoid freeze / thaw cycles.

Alternative Names:


MCM4 antibody; DNA replication licensing factor MCM4 antibody; MCM4 antibody; CDC21 antibody

SwissProt:

P33991

Product Description

The protein encoded by this gene is one of the highly conserved mini-chromosome maintenance proteins (MCM) that are essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by MCM proteins is a key component of the pre-replication complex (pre_RC) and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 6 and 7 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. The phosphorylation of this protein by CDC2 kinase reduces the DNA helicase activity and chromatin binding of the MCM complex. This gene is mapped to a region on the chromosome 8 head-to-head next to the PRKDC/DNA-PK, a DNA-activated protein kinase involved in the repair of DNA double-strand breaks. Alternatively spliced transcript variants encoding the same protein have been reported.



Western blot analysis of extracts of various cell lines, using MCM4 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!