

Anti Acetyl Histone Antibody

Catalog No: tcsa24027



Available Sizes

Size: 100µl



Specifications

Application:

WB, IHC, IF

Species Reactivity:

Human, Mouse, Rat

Host Species:

Rabbit

Immunogen / Amino acids:

A specific peptide of human Acetyl-Histone H2A-K9

Conjugation:

Unconjugated

Isotype:

IgG

Form:

Liquid

Storage Buffer:

PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Recommended Dilution:

WB 1:500 - 1:2000

IHC 1:500 - 1:1000

IF 1:100 - 1:500

Storage Instruction:


Store at -20C. Avoid freeze / thaw cycles.

Alternative Names:


HIST1H2AB antibody; Histone H2A type 1-B/E antibody; Histone H2A.2 antibody; Histone H2A/a antibody; Histone H2A/m antibody; HIST1H2AB antibody; H2AFM; antibody; HIST1H2AE antibody; H2AFA antibody

SwissProt:


P04908




Western blot analysis of extracts of various cell lines, using Acetyl-Histone H2A-K9 antibody. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST.




Immunofluorescence analysis of A549 cells using Acetyl-Histone H2A-K9 antibody.




Immunofluorescence analysis of HeLa cells using Acetyl-Histone H2A-K9 antibody.




Immunohistochemistry of paraffin-embedded mouse colon using Acetyl-Histone H2A-K9 antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human mammary cancer using Acetyl-Histone H2A-K9 antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human tonsil using Acetyl-Histone H2A-K9 antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human colon carcinoma using Acetyl-Histone H2A-K9 antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human lung using Acetyl-Histone H2A-K9 antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded mouse thyroid using Acetyl-Histone H2A-K9 antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded mouse testis using Acetyl-Histone H2A-K9 antibody at dilution of 1:100 (40x lens).



Immunofluorescence analysis of MCF-7 cells using Acetyl-Histone H2A-K9 antibody.

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!