

Polyclonal Antibody to Ubiquitin Carboxyl Terminal Hydrolase L1 (UCHL1)

Catalog No: tcua1117



Available Sizes

Size: 20ul

Size: 100ul

Size: 200ul



Specifications

Application:

WB; IHC; ICC; IP.

Species Reactivity:

Homo sapiens (Human)

Host Species:

Rabbit

Immunogen / Amino acids:

Recombinant Ubiquitin Carboxyl Terminal Hydrolase L1 (UCHL1)

Conjugation:

Unconjugated

Clonality:

Polyclonal

Form:

Liquid

Storage Buffer:

0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

Recommended Dilution:

Western blotting: 0.5-5 μ g/mL

Immunohistochemistry: 5-30 μ g/mL

Immunocytochemistry: 5-30 μ g/mL

Optimal working dilutions must be determined by end user.

Alternative Names:

PARK5; PGP9.5; Uch-L1; Neuron cytoplasmic protein 9.5; Ubiquitin thioesterase L1; Ubiquitin carboxyl-terminal hydrolase isozyme L1




Figure. Western Blot; Sample: Lane1: Mouse Brain Tissue; Lane2: Porcine Brain Tissue; Lane3: Human U-87MG Cells; Lane4: Human A549 Cells.)




Figure. Western Blot; Sample: Recombinant UCHL1, Human.)




Figure:FITC staining on IHC-P
Simple: A549 cells)



Figure. DAB staining on IHC-P;
Samples: Human Glioma Tissue.)




Figure. DAB staining on IHC-P;
Samples: Human Brain Tissue.)




Figure. DAB staining on IHC-P;
Samples: Human Lung Cancer Tissue.)




Figure. DAB staining on IHC-P; Samples: Human Kidney Tissue.)

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!