

Succimer

Catalog No: tcsc0013797



Available Sizes

Size: 100mg



Specifications

CAS No:

304-55-2

Formula:

$C_4H_6O_4S_2$

Pathway:

Others

Target:

Others

Purity / Grade:

>98%

Solubility:

DMSO : 300 mg/mL (1646.36 mM; Need ultrasonic and warming)

Alternative Names:

Dimercaptosuccinic acid;DMSA

Observed Molecular Weight:


182.22

Product Description

Succimer is a widely used chelating agent for the treatment of Pb poisoning.

In Vivo: Succimer is a widely used chelating agent for the treatment of Pb poisoning. Red blood cells (RBCs) from lead exposed animals treated with NAC or Succimer are shown to have significantly higher glutathione (GSH) levels and diminished malondialdehyde (MDA) levels when compare to the lead group. Succimer administration also results in decreased glucose 6-

phosphate dehydrogenase (G6PD) activity in RBCs from lead exposed animals^[1]. Succimer treatment produces a significant reduction in blood lead levels for both lead exposure conditions: the High Pb-succ group has blood lead levels that are 27% of the blood lead levels of the High Pb group at the end of treatment. Succimer is effective in substantially reducing brain lead, as brain lead levels in the High Pb-succ group are 37% of levels in the High Pb group^[2].



All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!