



## Pictilisib (dimethanesulfonate)

**Catalog No: tcsc0082** 

Available Sizes
Size: 10mg
Size: 50mg
Size: 100mg
Size: 200mg
Size: 500mg
Size: 1g
Size: 2g
Specifications
<b>CAS No:</b> 957054-33-0
Formula: ${\rm C_{25}^{\rm H}_{35}^{\rm N}_7^{\rm O}_9^{\rm S}_4^{\rm A}}$
Pathway: PI3K/Akt/mTOR;Autophagy
Target: PI3K;Autophagy
Purity / Grade: >98%
Solubility: 10 mM in DMSO





## **Alternative Names:**

GDC-0941 (dimethanesulfonate); GDC-0941 (2 MeSO3H salt)

## **Observed Molecular Weight:**

705.85

## **Product Description**

Pictilisib (GDC-0941) dimethanesulfonate is a potent inhibitor of **PI3K** $\alpha/\delta$  with **IC**<sub>50</sub> of 3 nM, with modest selectivity against p110 $\beta$  (11-fold) and p110 $\gamma$  (25-fold).

IC50 & Target: IC50: 3 nM (PI3K $\alpha$ ), 3 nM (PI3K $\delta$ )<sup>[5]</sup>

*In Vitro:* Pictilisib (GDC-0941) and docetaxel reduce tumor cell viability by 80% or greater in the breast cancer cell lines than single-agent treatment. GDC-0941 inhibits Akt phosphorylation and downstream targets of Akt signaling such as pPRAS40 and pS6 in Hs578T1.2 (PI3Kα wild-type), MCF7-neo/HER2 (PI3Kα-mutant), and MX-1 (PTEN-null) tumor models. Pictilisib (GDC-0941) decreases the time of docetaxel-induced mitotic arrest prior to apoptosis<sup>[1]</sup>. Pictilisib (GDC-0941) shows a high efficacy of antitumor activity in two gefitinib-resistant non-small cell lung cancer (NSCLC) cell lines, A549 and H460. Pictilisib (GDC-0941) is highly efficacious in combination with U0126 in inducing cell growth inhibition, G0-G1 arrest and cell apoptosis. H460 cells with activating mutations of PIK3CA are relatively more sensitive to Pictilisib (GDC-0941) than A549 cells with wild-type PIK3CA<sup>[3]</sup>. Pictilisib (GDC-0941) reduces PI3K pathway activity in both cell lines, illustrated by decreased pAK. Pictilisib (GDC-0941) significantly reduces secreted VEGF detected in the medium after hypoxic/anoxic exposure in all cells<sup>[4]</sup>.

*In Vivo:* Pictilisib (GDC-0941) (150 mg/kg, p.o.) leads to tumor stasis in MCF7-neo/HER2-bearing animals model. Pictilisib (GDC-0941) and docetaxel result in tumor regressions during the treatment period leading to enhanced antitumor responses<sup>[1]</sup>. Tumours in the Pictilisib (GDC-0941)-treated mice show a marked non-linear shrinkage, and when the Pictilisib (GDC-0941) treatment ceased, the tumours in the test cohort mice grow again<sup>[2]</sup>. GDC-0941Pictilisib (GDC-0941) (25 or 50 mg/kg) reduces tumor growth and PI3K and HIF-1 pathway activity in eGFP-FTC133 tumor-bearing mice<sup>[4]</sup>.

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!