



UK-5099

Catalog No: tcsc1048

| Available Sizes |
|---|
| Size: 5mg |
| Size: 10mg |
| Size: 50mg |
| Size: 100mg |
| Specifications |
| CAS No: 56396-35-1 |
| Formula: C ₁₈ H ₁₂ N ₂ O ₂ |
| Pathway: Membrane Transporter/Ion Channel |
| Target: Monocarboxylate Transporter |
| Purity / Grade: >98% |
| Solubility: H2O: |
| Alternative Names: PF-1005023 |
| Observed Molecular Weight: 288.3 |





Product Description

UK-5099 is a potent inhibitor of the mitochondrial pyruvate carrier (**MPC**). UK-5099 ihibits pyruvate-dependent O_2 consumption with an IC_{50} of 50 nM.

IC50 & Target: IC50: 50 nM (MPC)[1]

In Vitro: The trypanosomal pyruvate carrier is found to be rather insensitive to inhibition by alpha-cyano-4-hydroxycinnamate ($K_i = 17 \text{ mM}$) but can be completely blocked by UK-5099 ($K_i = 49 \text{ microM}$)^[2]. UK-5099 also inhibits the monocarboxylate transporter (MCT) ^[3]. UK5099 significantly inhibits the glucose-stimulated rise in oxygen consumption in a dose-dependent manner and at 150 μ M reduced oxygen consumption below basal levels. UK5099 reduces ATP levels and increases ADP and AMP levels in 832/13 cells^[4]. The UK5099 treated cells show significantly higher proportion of side population fraction and express higher levels of stemness markers Oct3/4 and Nanog. UK5099 application may be an ideal model for Warburg effect studies^[5].

In Vivo: The MPC inhibitor UK5099 increases the glucose excursion seen during an intraperitoneal glucose tolerance test in C57BLK mice^[4].

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!