



Reparixin

Catalog No: tcsc1379

Available Sizes
Size: 2mg
Size: 5mg
Size: 10mg
Size: 25mg
Size: 50mg
Size: 100mg
Size: 200mg
Specifications
CAS No: 266359-83-5
Formula: C ₁₄ H ₂₁ NO ₃ S
Pathway: GPCR/G Protein;Immunology/Inflammation
Target: CXCR;CXCR
Purity / Grade: >98%
Solubility: DMSO : ≥ 500 mg/mL (1764.35 mM)





Alternative Names:

Repertaxin; DF 1681Y

Observed Molecular Weight:

283.39

Product Description

Reparixin is a potent inhibitor of both CXCL8 receptors CXCR1/2, it inhibits weakly CXCR2-mediated cell migration (IC_{50} =100 nM), whereas it strongly blocks CXCR1-mediated chemotaxis (IC_{50} =1 nM).

IC50 & Target: IC50: 5.6/80 nM (CXCR1^{wt}/CXCR1^{lle43Val}, in L1.2 cell)^[1]

In Vitro: Reparixin is a potent functional inhibitor of CXCL8-induced biological activities on human PMNs with a marked selectivity (around 400-fold) for CXCR1, as shown in specific experiments on CXCR1/L1.2 and CXCR2/L1.2 transfected cells and on human PMNs. The efficacy of Reparixin is significantly lower in L1.2 cells expressing Ile43Val CXCR1 mutant (IC₅₀ values of 5.6 nM and 80 nM for CXCR1 wt and CXCR1 Ile43Val, respectively)^[1]. Reparixin is a non-competitive allosteric inhibitor of IL-8 receptors with a 400-fold higher efficacy in inhibiting CXCR1 activity than CXCR2^[2].

In Vivo: Reparixin is an inhibitor of CXCL8 receptor CXCR1 and CXCR2 activation, has been shown to attenuate inflammatory responses in various injury models. Spontaneously hypertensive rats (SHR) are administered a subcutaneous injection of Reparixin (5 mg/kg) daily for 3 weeks. Reparixin effectively decreases systolic blood pressure and increased the blood flow^[3]. Reparixin reduces the levels of IL-1 β in the brain after middle cerebral artery occlusion/reperfusion (MCAo) in mice. Bars represent levels of IL-1 β (pg/100 mg) measured by ELISA in the brain tissues of mice subjected or not (SHAM) to MCAo and pretreated with vehicle or Reparixin (30 mg/kg, s.c.)^[4].

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!