

# Tripelennamine (hydrochloride)

Catalog No: tcsc1395



## Available Sizes

**Size:** 100mg

**Size:** 200mg

**Size:** 500mg

**Size:** 1g

**Size:** 5g



## Specifications

**CAS No:**

154-69-8

**Formula:**

$C_{16}H_{22}ClN_3$

**Pathway:**

Immunology/Inflammation;GPCR/G Protein

**Target:**

Histamine Receptor;Histamine Receptor

**Purity / Grade:**

>98%

**Solubility:**

DMSO :  $\geq 33$  mg/mL (113.08 mM)

**Observed Molecular Weight:**

291.82

## Product Description

Tripelennamine HCl, a H<sub>1</sub>-receptor antagonist, is a psychoactive drug and member of the pyridine and ethylenediamine classes that is used as an antipruritic and first-generation antihistamine.


IC<sub>50</sub> Value:

Target: Histamine H<sub>1</sub> receptor

Tripelennamine can be used in the treatment of asthma, hay fever, rhinitis and urticaria.

in vitro: Arterial and mixed venous blood-gas and pH measurements were made at rest before and after saline or drug administration and during incremental exercise leading to maximal exertion at 14 m/s on 3.5% uphill grade for 120 s. Galloping at this workload elicited maximal heart rate and induced exercise-induced pulmonary hemorrhage in all horses in both treatments, thereby indicating that capillary stress failure-related pulmonary injury had occurred [1].

in vivo: The data obtained (median and range in brackets) in camels and horses, respectively, were as follows: the terminal elimination half-lives were 2.39 (1.91-6.54) and 2.08 (1.31-5.65) h, total body clearances were 0.97 (0.82-1.42) and 0.84 (0.64-1.17) L/h/kg. The volumes of distribution at steady state were 2.87 (1.59-6.67) and 1.69 (1.18-3.50) L/kg, the volumes of the central compartment of the two compartment pharmacokinetic model were 1.75 (0.68-2.27) and 1.06 (0.91-2.20) L/kg [2]. After intramuscular administration of 50 or 100 mg tripelennamine, mean plasma concentrations at 30 minutes were 105 and 194 ng/ml, respectively, and mean plasma t<sub>1/2</sub> values were 2.9 and 4.4 hours, respectively [3].



All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!