



Dabigatran

Catalog No: tcsc1399

Available Sizes
Size: 5mg
Size: 10mg
Size: 50mg
Size: 100mg
Specifications
CAS No: 211914-51-1
Formula: C ₂₅ H ₂₅ N ₇ O ₃
Pathway: Metabolic Enzyme/Protease
Target: Thrombin
Purity / Grade: >98%
Solubility: H2O :
Alternative Names: BIBR 953;BIBR 953ZW
Observed Molecular Weight: 471.51





Product Description

Dabigatran(BIB-953; BIBR 953ZW) is a reversible and selective, direct thrombin inhibitor (DTI) with Ki value of 4.5 nM.

IC50 Value: 4.5 nM (Ki); 10 nM(Thrombin-induced platelet aggregation) [1]

Target: thrombin

in vitro: Dabigatran selectively and reversibly inhibited human thrombin(Ki: 4.5 nM) as well as thrombin-induced platelet aggregation (IC(50): 10 nM), while showing no inhibitory effect on other platelet-stimulating agents. Thrombin generation in platelet-poor plasma (PPP), measured as the endogenous thrombin potential (ETP) was inhibited concentration-dependently (IC(50): 0.56 microM). Dabigatran demonstrated concentration-dependent anticoagulant effects in various species in vitro, doubling the activated partial thromboplastin time (aPTT), prothrombin time (PT) and ecarin clotting time (ECT) in human PPP at concentrations of 0.23, 0.83 and 0.18 microM, respectively [1].

in vivo: Dabigatran prolonged the aPTT dose-dependently after intravenous administration in rats (0.3, 1 and 3 mg/kg) and rhesus monkeys (0.15, 0.3 and 0.6 mg/kg). Dose- and time-dependent anticoagulant effects were observed with dabigatran etexilate administered orally to conscious rats (10, 20 and 50 mg/kg) or rhesus monkeys (1, 2.5 or 5 mg/kg), with maximum effects observed between 30 and 120 min after administration, respectively [1]. Patients treated with dabigatran etexilate experienced fewer ischaemic strokes (3.74 dabigatran etexilate vs 3.97 warfarin) and fewer combined intracranial haemorrhages and haemorrhagic strokes (0.43 dabigatran etexilate vs 0.99 warfarin) per 100 patient-years [2].

Clinical trial: An Evaluation of the Pharmacokinetics and Pharmacodynamics of Oral Dabigatran Etexilate in Hemodialysis Patients . Phase1

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!