

Batimastat (sodium salt)

Catalog No: tcsc1656



Available Sizes

Size: 1mg

Size: 5mg

Size: 10mg



Specifications

CAS No:

130464-84-5

Formula:

$C_{23}H_{30}N_3NaO_4S_2$

Pathway:

Metabolic Enzyme/Protease

Target:

MMP

Purity / Grade:

>98%

Solubility:

10 mM in DMSO

Alternative Names:

BB-94 sodium salt

Observed Molecular Weight:


499.62

Product Description

IC50 & Target: IC50: 3 nM (MMP-1), 4 nM (MMP-2), 4 nM (MMP-9), 6 nM (MMP-7), 20 nM (MMP-3)^[1]

In Vitro: Batimastat (BB-94) is a potent matrix metalloproteinase inhibitor, exhibits an unexpected mode of binding. Batimastat inhibits gelatinases A and B with IC₅₀ values of 4 nM and 10 nM, respectively. The IC₅₀ with the structurally similar collagenase Ht-d is 6 nM, which is comparable with values for MMP-1 (3 nM), MMP-8 (10 nM), and MMP-3 (20 nM)^[2]. CD30 shedding from the cell line Karpas299 can effectively be blocked by the hydroxamic acidbased metalloproteinase inhibitor Batimastat (BB-94, IC₅₀=230 nM)^[3].

In Vivo: Intraperitoneal administration of Batimastat (BB-94) effectively blocks growth of human ovarian carcinoma xenografts and murine melanoma metastasis and delays the growth of primary tumors in an orthotopic model of human breast cancer without cytotoxicity and without affecting mRNA levels^[2]. Batimastat (BB-94) is a synthetic matrix metalloproteinase inhibitor that has shown antineoplastic and antiangiogenic activity in various tumor models. Treatment with Batimastat (60 mg/kg i.p. every other day, for a total of eight injections) concomitantly with Cisplatin (4 mg/kg i.v., every 7 days for a total of three injections) completely prevents growth and spread of both xenografts, and all animals are alive and healthy on day 200^[4]. Kaplan-Meier analysis of survival (at 48 h) shows that animals treated with Batimastat (BB-94) have increased survival (95.2%) in comparison with controls (75%), and differences are almost statistically significant ($p=0.064$)^[5]. Matrix density is analyzed in saline- or Batimastat (40 mg/kg)-pretreated animals 4 h after E₂ administration, the time point at which collagen density is observed to be at its lowest after hormone treatment^[6].



All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!