

# **Benazepril (hydrochloride)**

## **Catalog No: tcsc1796**

Available Sizes

**Size:** 1g

Size: 5g

Specifications

CAS No:

86541-74-4

#### Formula:

C24H29CIN2O2

**Pathway:** Metabolic Enzyme/Protease

#### **Target:**

Angiotensin-converting Enzyme (ACE)

#### **Purity / Grade:**

>98%

## Solubility: H2O : 15.33 mg/mL (33.26 mM; Need ultrasonic); DMSO : $\geq$ 92 mg/mL (199.59 mM)

### Alternative Names:

CGS14824A

**Observed Molecular Weight:** 

460.95

## **Product Description**

Benazepril hydrochloride, an angiotensin converting enzyme inhibitor, which is a medication used to treat high blood pressure.

Copyright 2021 Taiclone Biotech Corp.



Target: angiotensin converting enzyme (ACE)

Benazepril hydrochloride is a medication used to treat high blood pressure (hypertension), congestive heart failure, and chronic renal failure. Upon cleavage of its ester group by the liver, benazepril hydrochloride is converted into its active form benazeprilat, a non-sulfhydryl angiotensin-converting enzyme (ACE) inhibitor [1].

Animals were randomly divided into 4 groups: sham STNx group (control), STNx group, morning benazepril hydrochloride group (MB) and evening benazepril hydrochloride group (EB).Benazepril hydrochloride was intragastrically administered at a dose of 10 mg/kg/day at 07:00 and 19:00 in the MB group and EB group respectively for 12 weeks. All the animals were synchronized to the light:dark cycle of 12:12 for 12 weeks. Systolic blood pressure (SBP), 24-h urinary protein excretion and renal function were measured at 11 weeks. Blood samples and kidneys were collected every 4 h throughout a day to detect the expression pattern of renin activity (RA), angiotensin II (AngII) and aldosterone (Ald) by radioimmunoassay (RIA) and the mRNA expression profile of clock genes (bmal1, dbp and per2) by real-time PCR at 12 weeks. Our results showed that no significant differences were noted in the SBP, 24-h urine protein excretion and renal function between the MB and EB groups. There were no significant differences in average Ald and RA content of a day between the MB group and EB group. The expression peak of bmal1 mRNA was phase-delayed by 4 to 8 h, and the diurnal variation of per2 and dbp mRNA diminished in the MB and EB groups compared with the control and STNx groups. It was concluded when the similar SBP reduction, RAAS inhibition and clock gene profile were achieved with optimal dose of benazepril hydrochloride, morning versus evening dosing of benazepril hydrochloride has the same renoprotection effects [2].

Clinical indications: Congestive heart failure; End stage renal disease; Hypertension

FDA Approved Date:

Toxicity: headaches; cough; Anaphylaxis; angioedema; hyperkalemia



#### All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!

Copyright 2021 Taiclone Biotech Corp.