

Etomidate

Catalog No: tcsc1833



Available Sizes

Size: 10mg

Size: 50mg

Size: 100mg

Size: 200mg

Size: 500mg



Specifications

CAS No:

33125-97-2

Formula:

$C_{14}H_{16}N_2O_2$

Pathway:

Neuronal Signaling;Membrane Transporter/Ion Channel

Target:

GABA Receptor;GABA Receptor

Purity / Grade:

>98%

Solubility:

DMSO : ≥ 100 mg/mL (409.35 mM)

Alternative Names:

R 16659

Observed Molecular Weight:

244.29

Product Description

Etomidate(R-16659) is a GABAA receptors agonist, which is a short acting intravenous anaesthetic agent used for the induction of general anaesthesia.


Target: GABA Receptor

Etomidate is a potent inhibitor of the adrenal response to surgery. The absence of clinical consequences associated with the blunted response suggests that a major increase in adrenal hormone production may not be necessary during surgery [1]. Etomidate is an intravenous induction agent that is associated with hemodynamic stability during intubation. The agent is therefore attractive for use in critically ill patients who have a high risk of hemodynamic instability during this procedure [2]. Etomidate use was not associated with all cause 28-day mortality or hospital mortality but was associated with significantly higher ICU mortality (91% vs. 64% for etomidate and controls groups, respectively; $p = 0.02$). Etomidate patients who received subsequent doses of hydrocortisone required lower doses of vasopressors and had more vasopressor-free days but no improvement in mortality [3].

Clinical indications:

FDA Approved Date: 1983

Toxicity: Undesirable side effects of etomidate that may limit its use include pain on injection, myoclonus and adrenocortical suppression lasting 4-6 hours following an induction dose.



All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!