

Tenovin-6

Catalog No: tcsc1976

Available Sizes

Size: 2mg

Size: 5mg

size: 10mg

Size: 100mg

Directifications

CAS No:

1011557-82-6

Formula:

 $C_{25}H_{34}N_4O_2S$

Pathway:

Autophagy; Epigenetics; Cell Cycle/DNA Damage; Apoptosis; Epigenetics; Cell Cycle/DNA Damage

Target:

Autophagy;Sirtuin;Sirtuin;MDM-2/p53;HDAC;HDAC

Purity / Grade:

>98%

Solubility:

DMSO : ≥ 31 mg/mL (68.19 mM)

Observed Molecular Weight:

454.63

Copyright 2021 Taiclone Biotech Corp.



Product Description

Tenovin-6 is a water soluble inhibitor of **SIRT1** and **SIRT2**, slightly inhibits **HDAC8**, and is also a potent activator of **p53**, with **IC**₅₀ s of 21 μ M, 10 μ M, and 67 μ M for SirT1, SirT2, and SirT3, respectively.

IC50 & Target: IC50: 21 μM (SirT1), 10 μM (SirT2), 67 μM (SirT3)^[1]

In Vitro: Tenovin-6 inhibits the growth of *S. cerevisiae* cultures with an IC_{50} of 30 µM and is more toxic to yeast than the less watersoluble tenovin-1. Tenovin-6 rapidly increases the levels of endogenous K382-Ac p53 in MCF-7 cells^[1]. Tenovin-6 (0 to 15 µM) dose dependently increases the level of LC3-II in diverse cell types, and the increase is ATG5/7 dependent. Tenovin-6 treatment also increases the number and intensity of autophagic vesicles with or without the presence of Torin 1, and prevents Torin 1-induced SQSTM1/p62 degradation. Tenovin-6 affects the acidification of autolysosomes and impairs the hydrolytic activity of lysosomes but does not affect the fusion between autophagosomes and lysosomes. That tenovin-6 inhibits autophagy does not correlate with p53 activation and SIRT1/2 inhibition by knockdown or knockout cannot mimic the effect of tenovin-6 on LC3B accumulation^[2]. Tenovin-6 (0, 1, 2.5, 5 or 10 µM) potently inhibits cell proliferation in a dose- and time-dependent manner in all OCI-Ly1, DHL-10, U2932, RIVA, HBL1 and OCI-Ly10 cell lines. Tenovin-6 consistently increases LC3B-II level in DLBCL cell lines by inhibiting the classical autophagy pathway, without activating p53, and the increase is independent of SIRT1/2/3 and p53. Tenovin-6 induces apoptosis through the extrinsic cell-death pathway^[3]. Tenovin-6 suppresses the growth of UM cells with IC₅₀ of 12.8 µM, 11.0 µM, 14.58 µM and 9.62 µM for 92.1, Mel 270, Omm 1 and Omm 2.3 cells, respectively^[4].

In Vivo: Tenovin-6 (50 mg/kg, i.p.) inhibits the growth of tumor in mice^[1].



All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!

Copyright 2021 Taiclone Biotech Corp.