

IWR-1 Catalog No: tcsc3107

Available Sizes

Size: 5mg

Size: 10mg

Size: 50mg

Specifications

CAS No:

1127442-82-3

Formula:

 $C_{25}H_{19}N_{3}O_{3}$

Pathway:

Stem Cell/Wnt

Target:

Wnt

Purity / Grade:

Solubility: DMSO : \geq 46 mg/mL (112.35 mM)

Alternative Names:

endo-IWR 1;IWR-1-endo

Observed Molecular Weight:

409.44

Product Description

Copyright 2021 Taiclone Biotech Corp.



IWR-1 is a **Wnt** pathway inhibitor with **IC**₅₀ of 180 nM in L-cells expressing Wnt3A, and induces Axin2 protein levels and promotes β -catenin phosphorylation by stabilizing Axin-scaffolded destruction complexes.

IC50 & Target: IC50: 180 nM (Wnt)

In Vitro: Both IWR-1 and XAV939 act as reversible Wnt pathway inhibitors and exhibit similar pharmacological effects in vitro. IWR-1 exerts its effect via interaction with Axin, while XAV939 binds TNKS directly^[1]. IWR-1 (10 μ M) induces stabilization of β -catenin disruption complex. IWR-1 (10 μ M) is added to the medium together with MIF, the size of cell colonies is extremely decreased, and that indicates the promoting effect of MIF on NSPC proliferation is inhibited by IWR-1 in any MIF concentration group. 2, 5 and 10 μ M of IWR-1 significantly inhibits the proliferation of NSPC dose-dependently. IWR-1 inhibites the promoting effect of MIF on NSPC dose-dependently. IWR-1 inhibites the promoting effect of MIF on NSPC dose-dependently. IWR-1 inhibites the promoting effect of MIF on NSPC dose-dependently. IWR-1 inhibites the promoting effect of MIF on NSPC dose-dependently. IWR-1 inhibites the promoting effect of MIF on NSPC dose-dependently. IWR-1 inhibites the promoting effect of MIF on NSPC dose-dependently. IWR-1 inhibites the promoting effect of MIF on NSPC dose-dependently. IWR-1 inhibites the promoting effect of MIF on NSPC dose-dependently. IWR-1 inhibites the promoting effect of MIF on NSPC dose-dependently. IWR-1 inhibites the promoting effect of MIF on NSPC dose-dependently. IWR-1 inhibites the promoting effect of MIF on NSPC dose-dependently. IWR-1 inhibites the promoting effect of MIF on NSPC dose-dependently. IWR-1 inhibites the promoting effect of IWR-1 (1 μ M). IWR-1 treatment partially reverses the FSH-induced inhibition of granulosa cell CARTPT mRNA expression^[3].



Copyright 2021 Taiclone Biotech Corp.