

SBE13 (Hydrochloride)

Catalog No: tcsc3785



Available Sizes

Size: 10mg

Size: 50mg



Specifications

CAS No:

1052532-15-6

Formula:

$C_{24}H_{28}Cl_2N_2O_4$

Pathway:

Cell Cycle/DNA Damage;Autophagy

Target:

Polo-like Kinase (PLK);Autophagy

Purity / Grade:

>98%

Solubility:

DMSO : ≥ 42 mg/mL (87.61 mM)

Observed Molecular Weight:

479.4


Product Description

SBE13 Hydrochloride is a potent and selective **Plk1** inhibitor, with an **IC₅₀** of 200 pM; SBE13 Hydrochloride poorly inhibits Plk2 (IC₅₀ >66 μ M) or Plk3 (IC₅₀=875 nM).

IC50 & Target: IC50: 200 pM (Plk1), >66 μ M (Plk2), 875 nM (Plk3)^[1]

In Vitro: SBE13 significantly reduce cell proliferation and induce apoptosis in HeLa cells, with an EC₅₀ of 18 μ M^[1]. SBE13 (1-100 μ M)

shows no effect on Caspase 3/7 activity in NIH-3T3 cells. SBE13 (66 and 100 μM) does not change morphology after treatment of primary cells. SBE13 (10 and 100 μM) reduces pRb staining in primary cells, and this indicates a G0/G1 arrest^[2]. SBE13 (66 and 100 μM) increases levels of cyclin B1, phospho histone H3, Wee1, Emi1 and securin, and results in cleavage of Cdc27 in HeLa cells. SBE13 (10 and 100 μM) also induces apoptosis of HeLa cells^[3].



All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!