



# **ELISA Kit for Trypsinogen Activation Peptide (TAP)**

**Catalog No: tcue427** 



## **Available Sizes**

Size: 96T



# **Specifications**

#### **Research Area:**

Infection immunity;Immune molecule;Hepatology;

#### **Species Reactivity:**

Mus musculus (Mouse)

### **Sample Type:**

Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids

## **Sensitivity:**

The minimum detectable dose of this kit is typically less than 46.4pg/mL

# **Detection Range:**

123.5-10,000pg/mL

#### **Assay Time:**

2h

#### **Detection Method:**

Enzyme-linked immunosorbent assay for Antigen Detection.

## **Tested Application:**

ELISA


# **Test Principle**

This assay employs the competitive inhibition enzyme immunoassay technique. A monoclonal antibody specific to Trypsinogen Activation Peptide (TAP) has been pre-coated onto a microplate. A competitive inhibition reaction is launched between biotin labeled Trypsinogen Activation Peptide (TAP) and unlabeled Trypsinogen Activation Peptide (TAP) (Standards or samples) with the pre-coated antibody specific to Trypsinogen Activation Peptide (TAP). After incubation the unbound conjugate is washed off. Next, avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. The amount of bound HRP conjugate is





reverse proportional to the concentration of Trypsinogen Activation Peptide (TAP) in the sample. After addition of the substrate solution, the intensity of color developed is reverse proportional to the concentration of Trypsinogen Activation Peptide (TAP) in the sample.



All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!