



## **ELISA Kit for Mucosal Addressin Cell Adhesion Molecule 1 (MAdCAM1)**

**Catalog No: tcue836** 

Q13477


| Available Sizes  |
|--|
| Size: 96T  |
| Specifications   |
| Research Area: CD & Adhesion molecule;   |
| Species Reactivity:<br>Homo sapiens (Human)  |
| Sample Type:<br>Serum, plasma, tissue homogenates, urine and other biological fluids |
| Sensitivity: The minimum detectable dose of this kit is typically less than 37pg/mL  |
| <b>Detection Range:</b> 78-5,000pg/mL  |
| Assay Time:<br>3h  |
| <b>Detection Method:</b> Enzyme-linked immunosorbent assay for Antigen Detection.    |
| Tested Application:<br>ELISA   |
| SwissProt:   |





## **Test Principle**

The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Mucosal Addressin Cell Adhesion Molecule 1 (MAdCAM1). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Mucosal Addressin Cell Adhesion Molecule 1 (MAdCAM1). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Mucosal Addressin Cell Adhesion Molecule 1 (MAdCAM1), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm  $\pm$  10nm. The concentration of Mucosal Addressin Cell Adhesion Molecule 1 (MAdCAM1) in the samples is then determined by comparing the O.D. of the samples to the standard curve.



All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!