

# ELISA Kit for Squamous Cell Carcinoma Antigen 2 (SCCA2)

Catalog No: tcue2110

Available Sizes

**Size:** 96T



**Research Area:** Tumor immunity;Gastroenterology;

## **Species Reactivity:**

Mus musculus (Mouse)

#### Sample Type:

Tissue homogenates, cell lysates and other biological fluids.

## Sensitivity:

The minimum detectable dose of this kit is typically less than 0.056ng/mL

#### **Detection Range:**

0.156-10ng/mL

## **Assay Time:** 3h

### **Detection Method:**

Enzyme-linked immunosorbent assay for Antigen Detection.

## **Tested Application:**

ELISA

## SwissProt:


Q9Z2G2

Copyright 2021 Taiclone Biotech Corp.



## **Test Principle**

The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Squamous Cell Carcinoma Antigen 2 (SCCA2). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Squamous Cell Carcinoma Antigen 2 (SCCA2). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Squamous Cell Carcinoma Antigen 2 (SCCA2), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm  $\pm 10$ nm. The concentration of Squamous Cell Carcinoma Antigen 2 (SCCA2) in the samples is then determined by comparing the O.D. of the samples to the standard curve.



All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!

Copyright 2021 Taiclone Biotech Corp.