

Polyclonal Antibody to Tumor Necrosis Factor Receptor Superfamily, Member 1A (TNFRSF1A)

Catalog No: tcua4807



Available Sizes

Size: 20ul

Size: 100ul

Size: 200ul



Specifications

Application:

WB; IHC; ICC; IP.

Species Reactivity:

Homo sapiens (Human)

Host Species:

Rabbit

Immunogen / Amino acids:

Recombinant Tumor Necrosis Factor Receptor Superfamily, Member 1A (TNFRSF1A)

Conjugation:

Unconjugated

Clonality:

Polyclonal

Form:

Liquid

Storage Buffer:

0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

Recommended Dilution:

Western blotting: 0.5-5 μ g/mL

Immunocytochemistry in formalin fixed cells: 5-20 μ g/mL

Immunohistochemistry in formalin fixed frozen section: 5-20 μ g/mL

Immunohistochemistry in paraffin section: 5-20 μ g/mL

Optimal workin

Alternative Names:

CD120A; P55; TBP1; FPF; TNF-R; TNF-R-I; TNF-R55; TNFAR; TNFR1; TNFR55; TNFR60; P55-R; P60; Tumor necrosis factor receptor 1; Tumor necrosis factor-binding protein 1





Figure. Western Blot; Sample: Lane1: Human HeLa Cells; Lane2: Human Jurkat Cells.)



Knockout Varification: Lane 1: Wild-type HeLa cell lysate; Lane 2: TNFRSF1A knockout HeLa cell lysate; Predicted MW: 50,38,25kDa Observed MW: 23kDa Primary Ab: 3 μ g/ml Rabbit Anti-Human TNFRSF1A Antibody Second Ab: 0.2 μ g/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody Catalog: SAA544Rb19)




Figure. Western Blot; Sample: Recombinant TNFRSF1A, Human.)



Figure:FITC staining on IHC-P
Simple: HeLa cells)




Figure. DAB staining on IHC-P; Samples: Human Brain Tissue.)




Figure. DAB staining on IHC-P; Samples: Human Kidney Tissue.)




Figure. DAB staining on IHC-P; Samples: Human Lung Cancer Tissue.)

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!