

Polyclonal Antibody to Aspartate Aminotransferase (AST)

Catalog No: tcua5347

KO VALIDATED



Available Sizes

Size: 20ul

Size: 100ul

Size: 200ul



Specifications

Application:

WB; IHC; ICC; IP.

Species Reactivity:

Rattus norvegicus (Rat)

Host Species:

Rabbit

Immunogen / Amino acids:

Recombinant Aspartate Aminotransferase (AST)

Conjugation:

Unconjugated

Clonality:

Polyclonal

Form:

Liquid

Storage Buffer:

0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

Recommended Dilution:

Western blotting: 0.5-2 μ g/mL

Immunohistochemistry: 5-20 μ g/mL

Immunocytochemistry: 5-20 μ g/mL

Optimal working dilutions must be determined by end user.

Alternative Names:

cCAT; AAT; ASAT; SGOT; GOT1; Cysteine transaminase, cytoplasmic; Transaminase A; Aspartate Transaminase 1, Cytoplasmic; Glutamic-Oxaloacetic Transaminase 1, Soluble




Figure. Western Blot; Sample: Lane1: Rat Brain Tissue; Lane2: Rat Heart Tissue; Lane3: Rat Liver Tissue; Lane4: Rat Kidney Tissue.)




Figure. Western Blot; Sample: Recombinant AST, Rat.)




Figure: FITC staining on IHC-P (Sample: HeLa cells)




Figure. DAB staining on IHC-P; Samples: Rat Liver Tissue.)




Figure. DAB staining on IHC-P;
Samples: Rat Testis Tissue.)




Figure. DAB staining on IHC-P;
Samples: Rat Kidney Tissue.)




Figure. DAB staining on IHC-P; Samples: Rat Intestine Tissue.)

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!