

Polyclonal Antibody to Nucleoporin 50kDa (NUP50)

Catalog No: tcua5501



Available Sizes

Size: 20ul

Size: 100ul

Size: 200ul



Specifications

Application:

WB; IHC; ICC; IP.

Species Reactivity:

Homo sapiens (Human)

Host Species:

Rabbit

Immunogen / Amino acids:

Recombinant Nucleoporin 50kDa (NUP50)

Conjugation:

Unconjugated

Clonality:

Polyclonal

Form:

Liquid

Storage Buffer:

0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

Recommended Dilution:

Western blotting: 0.5-2µg/ml

Immunocytochemistry in formalin fixed cells: 5-20 μ g/mL
Immunohistochemistry in formalin fixed frozen section: 5-20 μ g/mL
Immunohistochemistry in paraffin section: 5-20 μ g/mL
Optimal workin

Alternative Names:

NPAP60; NPAP60L; Nuclear pore-associated protein 60 kDa-like




Figure. Western Blot; Sample: Lane1: Human Jurkat Cells; Lane2: Human HeLa Cells; Lane3: Human HepG2 Cells.)




Figure. Western Blot; Sample: Recombinant NUP50, Human.)




Figure:FITC staining on IHC-P
Simple: HeLa cells)




Figure. DAB staining on IHC-P;
Samples: Human Skin Cancer Tissue.)




Figure. DAB staining on IHC-P;
Samples: Human Prostate Gland Cancer Tissue.)




Figure. DAB staining on IHC-P;
Samples: Human Cholangiocarcinoma Tissue.)




Figure. DAB staining on IHC-P; Samples: Human Brain Tissue.)

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!