

Acetyl-Histone H4-K5 mAb

Catalog No: tcba1486



Available Sizes

Size: 50ul

Size: 100ul

Size: 200ul



Specifications

Application:

WB,IHC,IF,IP

Species Reactivity:

Human,Mouse,Rat

Host Species:

Rabbit

Isotype:

IgG

Form:

Liquid

Storage Buffer:

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Recommended Dilution:

WB 1:500 - 1:2000

IHC 1:50 - 1:200

IF 1:50 - 1:200

IP 1:20 - 1:50

Storage Instruction:

Store at -20°C. Avoid freeze / thaw cycles.

Alternative Names:

H4/I;H4FI

SwissProt:

P62805

Gene ID:

8366 (human);

Calculated Molecular Weight:

11kDa

Purification:


Affinity purification

Cellular Location:

Chromosome,Nucleus,

Product Description

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H4 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6.




Western blot analysis of extracts of various cell lines, using Acetyl-Histone H4-K5 antibody.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.

Lysates/proteins: 25ug per lane.


Blocking buffer: 3% nonfat dry milk in TBST.




Immunohistochemistry of paraffin-embedded mouse brain using Acetyl-Histone H4-K5 antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human tonsil using Acetyl-Histone H4-K5 antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human colon carcinoma using Acetyl-Histone H4-K5 antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded mouse testis using Acetyl-Histone H4-K5 antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded mouse colon using Acetyl-Histone H4-K5 antibody at dilution of 1:100 (40x lens).



Immunofluorescence analysis of CRC cells using Acetyl-Histone H4-K5 antibody.



Immunofluorescence analysis of HeLa cells using Acetyl-Histone H4-K5 antibody.



Immunofluorescence analysis of SH-SY-5Y cells using Acetyl-Histone H4-K5 antibody.

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!