

Acetyl-Histone H2A-K9 mAb

Catalog No: tcba1489



Available Sizes

Size: 50ul

Size: 100ul

Size: 200ul



Specifications

Application:

WB,IHC,IF

Species Reactivity:

Human,Mouse,Rat

Host Species:

Rabbit

Isotype:

IgG

Form:

Liquid

Storage Buffer:

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Recommended Dilution:

WB 1:500 - 1:2000

IHC 1:500 - 1:1000

IF 1:100 - 1:500

Storage Instruction:

Store at -20°C. Avoid freeze / thaw cycles.

SwissProt:

P04908

Gene ID:


3012/8335 (human);

Calculated Molecular Weight:

14kDa

Purification:

Affinity purification




Western blot analysis of extracts of various cell lines, using Acetyl-Histone H2A-K9 antibody.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.

Lysates/proteins: 25ug per lane.


Blocking buffer: 3% nonfat dry milk in TBST.




Immunofluorescence analysis of A549 cells using Acetyl-Histone H2A-K9 antibody.




Immunofluorescence analysis of HeLa cells using Acetyl-Histone H2A-K9 antibody.




Immunohistochemistry of paraffin-embedded mouse colon using Acetyl-Histone H2A-K9 antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human mammary cancer using Acetyl-Histone H2A-K9 antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human tonsil using Acetyl-Histone H2A-K9 antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human colon carcinoma using Acetyl-Histone H2A-K9 antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human lung using Acetyl-Histone H2A-K9 antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded mouse thyroid using Acetyl-Histone H2A-K9 antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded mouse testis using Acetyl-Histone H2A-K9 antibody at dilution of 1:100 (40x lens).



Immunofluorescence analysis of MCF-7 cells using Acetyl-Histone H2A-K9 antibody.

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!