

Polyclonal Antibody to Prothrombin Fragment 1+2 (F1+2)

Catalog No: tcua6302



Available Sizes

Size: 20ul

Size: 100ul

Size: 200ul



Specifications

Application:

WB; IHC; ICC; IP.

Species Reactivity:

Homo sapiens (Human)

Host Species:

Rabbit

Immunogen / Amino acids:

Recombinant Prothrombin Fragment 1+2 (F1+2)

Conjugation:

Unconjugated

Clonality:

Polyclonal

Form:

Liquid

Storage Buffer:

0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

Recommended Dilution:

Western blotting: 0.5-2 μ g/mL

Immunohistochemistry: 5-20 μ g/mL

Immunocytochemistry: 5-20 μ g/mL

Optimal working dilutions must be determined by end user.




Figure. Western Blot; Sample: Lane1: Human 293T Cells; Lane2: Porcine Kidney Tissue; Lane3: Rat Serum.)




Figure. Western Blot; Sample: Recombinant F1+2, Human.)




Figure. DAB staining on IHC-P; Samples: Human Pancreas Cancer Tissue.)




Figure. DAB staining on IHC-P; Samples: Human Rectum Cancer Tissue.)




Figure. DAB staining on IHC-P; Samples: Human Rectum Cancer Tissue.)




Figure. DAB staining on IHC-P; Samples: Human Skin Cancer Tissue.)




Figure. DAB staining on IHC-P; Samples: Human Prostate Gland Cancer Tissue.)

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!