



## **HNRNPA1** Polyclonal Antibody

**Catalog No: tcba11158** 

IF 1:50 - 1:200

基	Available Sizes
Size:	50ul
Size:	100ul
Size:	200ul
	Specifications
<b>Appl</b> WB,IH	ication: IC,IF
	arch Area: Binding Protein(RBP),
	ies Reactivity: an,Mouse,Rat
<b>Host</b> Rabb	Species:
<b>Isoty</b> IgG	pe:
<b>Form</b>	
Storage Buffer: Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.	
WB 1	mmended Dilution: :500 - 1:2000 :50 - 1:200



Web: www.taiclone.com
Tel: +886-2-2735-9682
Email: order@taiclone.com

Storage Instruction: Store at -20°C. Avoid freeze / thaw cycles.
Alternative Names:
ALS19;ALS20;hnRNPA1;hnRNP-A1;HNRPA1L3;IBMPFD3;UP1
SwissProt: P09651
Gene ID: 3178 (human);
Calculated Molecular Weight: 29kDa/34kDa/38kDa
Purification: Affinity purification


## **Product Description**

**Cellular Location:** 

Cytoplasm, Nucleus,

This gene encodes a member of a family of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs), which are RNA-binding proteins that associate with pre-mRNAs in the nucleus and influence pre-mRNA processing, as well as other aspects of mRNA metabolism and transport. The protein encoded by this gene is one of the most abundant core proteins of hnRNP complexes and plays a key role in the regulation of alternative splicing. Mutations in this gene have been observed in individuals with amyotrophic lateral sclerosis 20. Multiple alternatively spliced transcript variants have been found. There are numerous pseudogenes of this gene distributed throughout the genome.





Western blot analysis of extracts of various cell lines, using HNRNPA1 antibody at 1:200 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.

Lysates/proteins: 25ug per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit.


Exposure time: 5min.




Immunohistochemistry of paraffin-embedded human brain astrocytoma using HNRNPA1 Antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human breast cancer using HNRNPA1 Antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded human stomach using HNRNPA1 Antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human leiomyoma of uterus using HNRNPA1 Antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human adenomyosis using HNRNPA1 Antibody at dilution of 1:100 (40x lens).







Immunohistochemistry of paraffin-embedded mouse liver using HNRNPA1 Antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded mouse kidney using HNRNPA1
Antibody at dilution of 1:100 (40x lens).




Immunofluorescence analysis of A549 cells using HNRNPA1 antibody.



Immunohistochemistry of paraffin-embedded human gastric injury using HNRNPA1 Antibody at dilution of 1:100 (40x lens).





Immunohistochemistry of paraffin-embedded human breast cancer using HNRNPA1 Antibody at dilution of 1:100 (40x lens).





Immunohistochemistry of paraffin-embedded human liver cancer using HNRNPA1 Antibody at dilution of 1:100 (40x lens).















dilution of 1:100 (40x lens).



All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!