

# Polyclonal Antibody to Proliferating Cell Nuclear Antigen (PCNA)

## Catalog No: tcua6581



### Available Sizes

---

**Size:** 20ul

---

**Size:** 100ul

---

**Size:** 200ul

---



### Specifications

---

**Application:**

WB; IHC; ICC; IP.

---

**Species Reactivity:**

Homo sapiens (Human), Mus musculus (Mouse), Rattus norvegicus (Rat), Oryctolagus cuniculus (Rabbit), Rhesus monkey (Simian), Canis familiaris; Canine (Dog), Sus scrofa; Porcine (Pig)

---

**Host Species:**

Rabbit

---

**Immunogen / Amino acids:**

Recombinant Proliferating Cell Nuclear Antigen (PCNA)

---

**Conjugation:**

Unconjugated

---

**Clonality:**

Polyclonal

---

**Form:**

Liquid

---

**Storage Buffer:**

0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

---

**Recommended Dilution:**

Western blotting: 0.5-2 $\mu$ g/mL;1:200-800

Immunohistochemistry: 5-20 $\mu$ g/mL;1:20-80

Immunocytochemistry: 5-20 $\mu$ g/mL;1:20-80

Optimal working dilutions must be determined by end user.

**Alternative Names:**

Cyclin




Figure. Western Blot; Sample: Lane1: Human Liver Tissue; Lane2: Human MCF7 Cells.)




Figure. Western Blot; Sample: Recombinant PCNA, Human.)




Figure. DAB staining on IHC-P; Samples: Human Liver Cancer Tissue.)



Figure. DAB staining on IHC-P; Samples: Human Stomach Tissue.)




Figure. DAB staining on IHC-P;  
Samples: Human Rectum Tissue.)




Figure. DAB staining on IHC-P; Samples: Human Kidney Tissue.)

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!