

Polyclonal Antibody to Cyclin D2 (CCND2)

Catalog No: tcua6588



Available Sizes

Size: 20ul

Size: 100ul

Size: 200ul



Specifications

Application:

WB; IHC; ICC; IP.

Species Reactivity:

Mus musculus (Mouse)

Host Species:

Rabbit

Immunogen / Amino acids:

Recombinant Cyclin D2 (CCND2)

Conjugation:

Unconjugated

Clonality:

Polyclonal

Form:

Liquid

Storage Buffer:

0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

Recommended Dilution:

Western blotting: 0.5-5μg/mL

Immunohistochemistry: 5-20μg/mL

Immunocytochemistry: 5-20μg/mL

Optimal working dilutions must be determined by end user.

Alternative Names:

CCN-D2; G1/S-Specific Cyclin-D2




Figure. Western Blot; Sample: Lane1: Human 293T Cells; Lane2: Human Hela Cells.)




Figure. Western Blot; Sample: Recombinant CCND2, Mouse.)




Figure. DAB staining on IHC-P;
Samples: Mouse Testis Tissue.)




Figure. DAB staining on IHC-P;
Samples: Mouse Kidney Tissue.)




Figure. DAB staining on IHC-P;
Samples: Mouse Stomach Tissue.)




Figure. DAB staining on IHC-P; Samples: Mouse Intestine Tissue.)

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!