

Polyclonal Antibody to Intercellular Adhesion Molecule 1 (ICAM1)

Catalog No: tcua6691



Available Sizes

Size: 20ul

Size: 100ul

Size: 200ul



Specifications

Application:

WB; IHC; ICC; IP.

Species Reactivity:

Rattus norvegicus (Rat)

Host Species:

Rabbit

Immunogen / Amino acids:

Recombinant Intercellular Adhesion Molecule 1 (ICAM1)

Conjugation:

Unconjugated

Clonality:

Polyclonal

Form:

Liquid

Storage Buffer:

0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

Recommended Dilution:

Western blotting: 0.5-5 μ g/mL

Immunohistochemistry: 5-20 μ g/mL

Immunocytochemistry: 5-20 μ g/mL

Optimal working dilutions must be determined by end user.

Alternative Names:

CD54; BB2; P3.58; Human Rhinovirus Receptor; Major group rhinovirus receptor




Figure. Western Blot; Sample: Lane1: Human HeLa Cells; Lane2: Human K562 Cells.)




Figure. Western Blot; Sample: Recombinant ICAM1, Rat.)




Figure:FITC staining on IHC-P
Simple: HeLa cells)




Figure. DAB staining on IHC-P;
Samples: Rat Lung Tissue.)




Figure. DAB staining on IHC-P;
Samples: Rat Liver Tissue.)




Figure. DAB staining on IHC-P;
Samples: Rat Spinal Cord Tissue.)




Figure. DAB staining on IHC-P; Samples: Rat Brain Tissue.)

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!