

Polyclonal Antibody to Acid Phosphatase 3, Prostatic (ACP3)

Catalog No: tcua7412



Available Sizes

Size: 20ul

Size: 100ul

Size: 200ul



Specifications

Application:

WB; IHC; ICC; IP.

Species Reactivity:

Homo sapiens (Human)

Host Species:

Rabbit

Immunogen / Amino acids:

Recombinant Acid Phosphatase 3, Prostatic (ACP3)

Conjugation:

Unconjugated

Clonality:

Polyclonal

Form:

Liquid

Storage Buffer:

0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

Recommended Dilution:

Western blotting: 0.5-5 μ g/mL

Immunohistochemistry: 5-20 μ g/mL

Immunocytochemistry: 5-20 μ g/mL

Optimal working dilutions must be determined by end user.

Alternative Names:

PSAP; ACP3; PAP; PAPf39; 5'-NT; Prostatic Specific Acid Phosphatase; Ecto-5'-Nucleotidase




Figure . Western Blot; Sample:Lane1: Human PC-3 Cells; Lane2: Human MCF7 Cells.)




Figure. Western Blot; Sample: Recombinant ACP3, Human.)




Figure:FITC staining on IHC-P
Simple:Mcf7 cells)




Figure. DAB staining on IHC-P;
Samples: Human Glioma Tissue.)




Figure. DAB staining on IHC-P;
Samples: Human Kidney Tissue.)




Figure. DAB staining on IHC-P;
Samples: Human Stomach Tissue.)




Figure. DAB staining on IHC-P; Samples: Human Liver Tissue.)

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!