

GAPDH Polyclonal Antibody

Catalog No: tcba13637



Available Sizes

Size: 50ul

Size: 100ul

Size: 200ul



Specifications

Application:

WB,IHC,IF

Research Area:

Epigenetics & Nuclear Signaling,Epigenetic Modifications,Epigenetic Modifications_Methylation,Epigenetics & Nuclear Signaling,Epigenetic Modifications,Epigenetic Modifications_Methylation,

Species Reactivity:

Human,Mouse,Rat

Host Species:

Rabbit

Immunogen / Amino acids:

Recombinant fusion protein containing a sequence corresponding to amino acids 1-335 of human GAPDH (NP_002037.2).

Conjugation:

Unconjugated

Clonality:

Polyclonal

Isotype:

IgG

Storage Buffer:

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Recommended Dilution:

WB 1:1000 - 1:10000

IHC 1:50 - 1:100

IF 1:20 - 1:50

Storage Instruction:

Store at -20°C. Avoid freeze / thaw cycles.

Alternative Names:

G3PD;GAPD;HEL-S-162eP

SwissProt:

P04406

Gene ID:

2597 (human);

Calculated Molecular Weight:

31kDa/36kDa

Purification:


Affinity purification

Cellular Location:


Cytoplasm,Membrane,Nucleus,cytoskeleton,cytosol,perinuclear region,

Product Description


This gene encodes a member of the glyceraldehyde-3-phosphate dehydrogenase protein family. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions. The product of this gene catalyzes an important energy-yielding step in carbohydrate metabolism, the reversible oxidative phosphorylation of glyceraldehyde-3-phosphate in the presence of inorganic phosphate and nicotinamide adenine dinucleotide (NAD). The encoded protein has additionally been identified to have uracil DNA glycosylase activity in the nucleus. Also, this protein contains a peptide that has antimicrobial activity against *E. coli*, *P. aeruginosa*, and *C. albicans*. Studies of a similar protein in mouse have assigned a variety of additional functions including nitrosylation of nuclear proteins, the regulation of mRNA stability, and acting as a transferrin receptor on the cell surface of macrophage. Many pseudogenes similar to this locus are present in the human genome. Alternative splicing results in multiple transcript variants.




Western blot analysis of extracts of various cell lines, using GAPDH antibody at 1:1000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.
Lysates/proteins: 25ug per lane.
Blocking buffer: 3% nonfat dry milk in TBST.




Immunohistochemistry of paraffin-embedded rat pancreas using GAPDH antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded human breast cancer using GAPDH antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded mouse heart using GAPDH antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded mouse heart using GAPDH antibody at dilution of 1:100 (40x lens).



Immunofluorescence analysis of MCF-7 cells using GAPDH antibody. Blue: DAPI for nuclear staining.

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!