

Goat Anti-Rabbit IgG (H+L) Rhodamine (TRITC) conjugated Antibody

Catalog No: tcba13716



Available Sizes

Size: 100ul



Specifications

Application:

IF,FC

Species Reactivity:

Rabbit

Host Species:

Goat

Immunogen / Amino acids:

Rabbit IgG

Conjugation:

TRITC

Clonality:

Polyclonal

Isotype:

TRITC conjugated IgG

Form:

Liquid

Storage Buffer:

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Recommended Dilution:

IHC 1:20 - 1:100

ICC 1:20 - 1:100

IF 1:20 - 1:100

FC 1:20 - 1:100

Amino Acid Sequence:


Affinity purification

Storage Instruction:

Store at -20°C. Avoid freeze / thaw cycles.

Reference Sequence No.:


Affinity purification



Immunofluorescence analysis of HeLa cells, using ACO2 antibody as the primary antibody at dilution of 1:100. The cells were incubated with the primary antibody overnight at 4°C.

Secondary antibody: Rhodamine (TRITC) Goat Anti-Rabbit IgG (H+L) at 1:200 dilution.

Blue: DAPI for nuclear staining.



Immunofluorescence analysis of HeLa cells, using PARP1 antibody as the primary antibody at dilution of 1:100. The cells were incubated with the primary antibody overnight at 4°C.

Secondary antibody: Rhodamine (TRITC) Goat Anti-Rabbit IgG (H+L) at 1:100 dilution.

Blue: DAPI for nuclear staining.

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!