

# Goat Anti-Rabbit IgG (H+L) Rhodamine (TRITC) conjugated Antibody

Catalog No: tcba13716



## Available Sizes

**Size:** 100ul



## Specifications

**Application:**

IF,FC

**Species Reactivity:**

Rabbit

**Host Species:**

Goat

**Immunogen / Amino acids:**

Rabbit IgG

**Conjugation:**

TRITC

**Clonality:**

Polyclonal

**Isotype:**

TRITC conjugated IgG

**Form:**

Liquid

**Storage Buffer:**

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Recommended Dilution:**

IHC 1:20 - 1:100

ICC 1:20 - 1:100

IF 1:20 - 1:100

FC 1:20 - 1:100

---

**Amino Acid Sequence:**

Affinity purification

---

**Storage Instruction:**


Store at -20°C. Avoid freeze / thaw cycles.

---

**Reference Sequence No.:**

Affinity purification


---



Immunofluorescence analysis of HeLa cells, using ACO2 antibody as the primary antibody at dilution of 1:100. The cells were incubated with the primary antibody overnight at 4°C.

Secondary antibody: Rhodamine (TRITC) Goat Anti-Rabbit IgG (H+L) at 1:200 dilution.

Blue: DAPI for nuclear staining.



Immunofluorescence analysis of HeLa cells, using PARP1 antibody as the primary antibody at dilution of 1:100. The cells were incubated with the primary antibody overnight at 4°C.

Secondary antibody: Rhodamine (TRITC) Goat Anti-Rabbit IgG (H+L) at 1:100 dilution.

Blue: DAPI for nuclear staining.

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!