



Mouse Anti-Human IgG Antibody- prediluted

Catalog No: tcna1545pre

Available Sizes
Size: 7ml
Specifications
Application: IHC-P
Species Reactivity: Human. Other species not known.
Host Species: Mouse
Immunogen / Amino acids: Human Ig Gamma Chain was used as the immunogen for this IgG antibody.
Conjugation: Unconjugated
Clonality: Monoclonal
Clones: GHC07-1
Isotype: Mouse IgG1, kappa
Form: Liquid
Storage Buffer: Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*



Recommended Dilution:

Prediluted IHC only format: incubate for 30 min at RT (2)The optimal dilution of the IgG antibody for each application should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer pH 6.0

for 10-20 min followed by cooling at RT for 20 minutes.

2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required)

drip mAb solution onto the tissue section and incubate at RT for 30 min.

Storage Instruction:

Store the IgG antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

SwissProt:


P01857, P01859, P01860, P01861

References

Protein G affinity chromatography

Product Description

Recognizes a protein of 75kDa, identified as gamma heavy chain of human immunoglobulins. It reacts with all sub-classes of gamma chain of human immunoglobulins. It does not cross-react with alpha (IgA), mu (IgM), epsilon (IgE), or delta (IgD), heavy chains, T-cells, monocytes, granulocytes, or erythrocytes. This mAb is useful in the identification of leukemias, plasmacytomas, and certain non-Hodgkin\'s lymphomas. The most common feature of these malignancies is the restricted expression of a single heavy chain class. Demonstration of clonality in lymphoid infiltrates indicates that the infiltrate is clonal and therefore malignant.



IHC testing of human tonsil and IgG antibody (GHC07-1)

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!