



Recombinant Goat Anti-Rabbit IgG Fc Antibody

Catalog No: tcna4754

Available Sizes
Size: 50ug
Size: 100ug
Specifications
Application: WB (non-reduced), ICC, IHC, FACS, ELISA
Species Reactivity: Rabbit
Host Species: Goat
Immunogen / Amino acids: Rabbit IgG was used as the immunogen for this recombinant Rabbit IgG Fc antibody.
Conjugation: Biotin Conjugate
Clonality: Recombinant Goat Monoclonal
Clones: RMG02
Isotype: Goat IgG
Form: Liquid
Storage Buffer:





1 mg/ml in PBS with 50% glycerol, 1% BSA and 0.09% sodium azide

Recommended Dilution:

ELISA: 0.01ug/ml - 0.5ug/ml

Immunocytochemistry: 0.5-2ug/ml

Immunohistochemistry (FFPE): 0.5-2ug/ml (1)

Western Blot (non-reduced): 0.1-0.5ug/mlThe stated application concentrations are suggested starting points. Titration of the recombinant Rabbit IgG Fc antibody may be required due to differences in protocols and secondary/substrate sensitivity.

1. A pH6 Citrate buffer or pH9 Tris/EDTA buffer HIER step is recommended for testing of FFPE tissue sections.

Storage Instruction:

Store the recombinant Rabbit IgG Fc antibody at -20oC (with glycerol) or aliquot and store at -20oC (without glycerol).

Gene ID:


N/A (human);


References

Biotin Conjugate

Product Description

This recombinant Rabbit IgG Fc antibody reacts to the Fc region of Rabbit IgG. No cross reactivity with human IgG, rat IgG, or mouse IgG.









ELISA Titration: the plate was coated with different amounts of rabbit IgG. A serial dilution of the recombinant Rabbit IgG Fc antibody was used as the primary and an alkaline phosphatase conjugated anti-goat IgG as the secondary.

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!