

# Mouse Anti-Human IgM Antibody

Catalog No: tcna537



## Available Sizes

---

**Size:** 20ug

**Size:** 100ug



## Specifications

---

**Application:**

FACS, IF, IHC-P

**Species Reactivity:**

Human. Other species not tested.

**Host Species:**

Mouse

**Immunogen / Amino acids:**

The Mu heavy chain of human Ig was used as the immunogen for the Anti-IgM antibody.

**Conjugation:**

Unconjugated

**Clonality:**

Monoclonal

**Clones:**

ICO-30

**Isotype:**

Mouse IgG1, kappa

**Form:**

Liquid

**Storage Buffer:**

0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide

---

**Concentration:**

0.2 mg/ml

---

**Recommended Dilution:**

Flow Cytometry: 0.5-1ug/million cells in 0.1ml

Immunofluorescence: 0.5-1ug/ml

Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT (1)

Prediluted format : incubate for 30 min at RT (2)Optimal dilution of the Anti-IgM antibody should be determined by the researcher.

1. Staining of formalin/paraffin tissues requires boiling tissue sections in 10mM Citrate buffer  
pH 6.0

for 10-20 min followed by cooling at RT for 20 min.

2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required)

drip mAb solution onto the tissue section and incubate at RT for 30 min.

---

**Storage Instruction:**

Store the Anti-IgM antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

---

**SwissProt:**

P01871 & P20769


---

**References**

Protein G affinity chromatography

**Product Description**

Recognizes a protein of 75kDa, identified as mu heavy chain of human immunoglobulins. It does not cross-react with alpha (IgA), gamma (IgG), epsilon (IgE), or delta (IgD), heavy chains, T-cells, monocytes, granulocytes, or erythrocytes. Monomeric IgM is expressed as a membrane bound antibody on the surface of B cells and as a pentamer when secreted by plasma cells. IgM antibody is prominent in early immune responses to most antigens. Aberrant levels are associated with immune deficiency states, hereditary deficiencies, myeloma, Waldenstrom's macroglobulinemia, chronic infection and hepatocellular disease. This mAb is useful in the identification of leukemias, plasmacytomas, and certain non-Hodgkins lymphomas. The most common feature of these malignancies is the restricted expression of a single heavy chain class. Demonstration of clonality in lymphoid infiltrates indicates that the infiltrate is clonal and therefore malignant.



Formalin-fixed, paraffin-embedded human tonsil stained with anti-IgM antibody (ICO-30)

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!