

Monoclonal Antibody to Kidney Injury Molecule 1 (Kim1)

Catalog No: tcua8853



Available Sizes

Size: 20ul

Size: 100ul

Size: 200ul



Specifications

Application:

WB; IHC; ICC; IP.

Species Reactivity:

Homo sapiens (Human)

Host Species:

Mouse

Immunogen / Amino acids:

Recombinant Kidney Injury Molecule 1 (Kim1)

Conjugation:

Unconjugated

Clonality:

Monoclonal

Clones:

C2

Isotype:

IgG2b Kappa

Form:

Liquid

Storage Buffer:

0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

Recommended Dilution:

Western blotting: 0.5-5 μ g/mL

Immunocytochemistry in formalin fixed cells: 5-30 μ g/mL

Immunohistochemistry in formalin fixed frozen section: 5-30 μ g/mL

Immunohistochemistry in paraffin section: 5-30 μ g/mL

Optimal working dilutions must be determined

Alternative Names:

HAVCR1; TIM1; TIMD1; HAVCR; Hepatitis A Virus Cellular Receptor 1; T Cell Immunoglobulin And Mucin Domain-Containing Protein 1; T-cell immunoglobulin mucin receptor 1




Figure. Western Blot; Sample: Recombinant Kim1, Human.)



Figure. DAB staining on IHC-P;
Samples: Human Liver Tissue.)




Figure. DAB staining on IHC-P;
Samples: Human Prostate Gland Tissue.)




Figure. DAB staining on IHC-P;
Samples: Human Stomach Tissue.)




Figure. DAB staining on IHC-P; Samples: Human Kidney Tissue.)

All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!