

# 293T / 17

Catalog No: tcel469



## Available Sizes

**Size:** 1×10<sup>6</sup>cells/t25culturebottle



## Specifications

### Subculturing:

Remove and discard culture medium. Briefly rinse the cell layer with DPBS solution to remove all traces of serum that contains trypsin inhibitor. Add 1.0 to 2.0 mL of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 2 to 3 minutes). Cells that are difficult to detach may be placed at 37°C to facilitate dispersal. Add 4.0 to 6.0 mL of complete growth medium and aspirate cells by gently pipetting. Add appropriate aliquots of the cell suspension to new culture vessels.

## Product Description

293T/17 cells were cotransfected with the pCRIPenv- and the pCRIPgag-2 vectors to obtain the ANJOU 65 cell line. ANJOU 65 cells were cotransfected with the pCRIPgag-2 and pGPT2E vectors to obtain the BOSC 23 ecotropic envelope-expression packaging cell line. ANJOU 65 cells were also cotransfected with the pCRIPAMgag vector along with a plasmid expressing the gpt resistance gene to obtain the Bing amphotropic envelope-expression packaging cell line.



All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!