

Rabbit anti-Human IgG Fc Antibody

Catalog No: tccua64832



Available Sizes

Size: 500ug



Specifications

Application:

ELISA

Species Reactivity:

Human

Host Species:

Rabbit

Immunogen / Amino acids:

Human IgG Fc fragment

Conjugation:

Non-conjugated

Isotype:

IgG

Form:

Liquid

Storage Buffer:

Preservative: 0.03% Proclin 300

Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

Storage Instruction:

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Product Description

The fragment crystallizable region (Fc region) is the tail region of an antibody that interacts with cell surface receptors called Fc receptors and some proteins of the complement system. This property allows antibodies to activate the immune system. In IgG, IgA and IgD antibody isotypes, the Fc region is composed of two identical protein fragments, derived from the second and third constant domains of the antibody two heavy chains; IgM and IgE Fc regions contain three heavy chain constant domains (CH domains 2-4) in each polypeptide chain. The Fc regions of IgGs bear a highly conserved N-glycosylation site. Glycosylation of the Fc fragment is essential for Fc receptor-mediated activity. The N-glycans attached to this site are predominantly core-fucosylated diantennary structures of the complex type. In addition, small amounts of these N-glycans also bear bisecting GlcNAc and α -2,6 linked sialic acid residues.



All products are for RESEARCH USE ONLY. Not for diagnostic & therapeutic purposes!